

General elastic interaction in nematic liquid crystals colloids

S. B. Chernyshuk ^{1,2}, and B. I. Lev ²,

¹, *Institute of Physics, NAS Ukraine, Prospekt Nauki 46, Kyiv 03650, Ukraine.* ², *Bogolyubov Institute of Theoretical Physics, NAS Ukraine, Metrologichna 14-b, Kyiv 03680, Ukraine.*

(Dated: July 29, 2009)

The new free energy functional that describes general elastic interaction between colloidal particles and nematic liquid crystal has been proposed. It generalizes results of the paper [7] on the case of arbitrary orientation of colloidal particles and is valid for arbitrary surface anchoring strength. Formal analogies and differences between electric particles and colloidal particles in LC are found. It is first time shown that spur of the quadrupole moment tensor is different from zero $Sp\hat{Q}_\mu \neq 0$ in the case of axial symmetry breaking of the director field around colloidal particle in contrast to electrostatics. New general presentation for the elastic interaction between arbitrary colloids is obtained which contains new multipole terms connected with $Sp\hat{Q}_\mu \neq 0$ that are absent in the standard electrostatics.

PACS numbers: 61.30.-v, 42.70.Df, 85.05.Gh

Colloidal particles in liquid crystals (LC) have attracted a great research interest during the last years. Anisotropic properties of the host fluid - liquid crystal give rise to a new class of colloidal anisotropic interactions that never occurs in isotropic hosts. Liquid crystal colloidal system have much recent interest as models for diverse phenomena in condense matter physics. The anisotropic interactions result in different structures of colloidal particles such as linear chains in inverted nematic emulsions [1, 2], 2D crystals [3] and 2D hexagonal structures at nematic-air interface [4, 5].

Study of anisotropic colloidal interactions has been made both experimentally [2]-[6] and theoretically [7]-[15]. The first theoretical approach was developed in [1, 7] with help of ansatz functions for the director and using multiple expansion in the far field area. Authors investigated spherical particles with hyperbolic hedgehog and found dipole and quadrupole elastic interactions between such particles. Another approach [8, 9] gave possibility to find approximate solutions in terms of geometrical shape of particles for the case of small anchoring strength and has provided the way to connect the type of the interaction potential with the local symmetry of the director field around particles [10]. The concept of coat has been introduced that contains all topological defects inside and carries the symmetry of the director and it enables to determine qualitatively the type of the interaction potential. It was first shown there that axial symmetry breaking leads to the origin of elastic charge and coulomb-like interaction. But the coat is not quantitatively exactly defined. In [11, 12] authors suggested approach by fixing director field on the surface of the imaginary sphere. There multiple expansion of the potential was obtained as well but coefficients differ significantly from the results of [7]. For instance dipole-dipole interaction was found to be three times weaker, and quadrupole five times weaker than in [7]. On the other hand authors of [6] recently have measured experimentally both interactions and found that results are in accordance with Lubensky at all prediction with about 10% accuracy. This allows to confirm rightness of the main assumptions of [7] for spherical particles though it do not work for particles with broken axial symmetry. In this paper we suggest to generalize that results for the case of arbitrary particle's orientation. We propose new free energy functional that describes interaction of colloidal particles with director field and turns into functional of [7] for axial particles. It is valid for both strong anchoring case with the presence of topological defects and for weak anchoring as well. We show that axial symmetry breaking leads to the violation of $Sp\hat{Q}_\mu = 0$ condition for spur of the quadrupole moment of particles in contrast to the classical electrostatics. This leads to the principal new multipole terms in elastic interaction connected with $Sp\hat{Q}_\mu$ which are absent in the classical electrostatics. We show that the reason of such difference leads in nonlinearities of director field on close distances and in surface driven effects associated with anchoring.

In order to explain subsequent speculations we consider first the case of homogeneous liquid crystal with uniform director $\mathbf{n}_0$ and one particle immersed into it. Anchoring of the liquid crystal with the surface of the particle produces deformations of the director field around the particle so that director $\mathbf{n}(\mathbf{R})$ varies from point to point. In the one-constant approximation the total free energy of the system: particle plus liquid crystal has the form:

$$F = \frac{K}{2} \int dV [(div\mathbf{n})^2 + (rot\mathbf{n})^2] + \oint dS W (\nu \cdot \mathbf{n})^2 \quad (1)$$

where W is anchoring strength coefficient, ν is the unit normal vector, integration $\oint dS$ is carried out on the surface of the particle, K is the Frank elastic constant.

Far from the particle director field variations are small $\mathbf{n}(\mathbf{R}) \approx (n_x, n_y, 1)$, $|n_\mu| \ll 1$ ($\mu = x, y$) and bulk free energy has the form:

$$F_{b,linear} = \frac{K}{2} \int dV \{(\nabla n_x)^2 + (\nabla n_y)^2\} \quad (2)$$

which brings Euler-Lagrange equation of Laplace type:

$$\Delta n_\mu = 0 \quad (3)$$

At large distances R in general case it can be expanded in multiples,

$$n_\mu(\mathbf{R}) = \frac{q_\mu}{R} + \frac{\mathbf{p}_\mu \mathbf{u}}{R^2} + 3 \frac{\mathbf{u} : \hat{Q}_\mu : \mathbf{u}}{R^3} - \frac{Sp\hat{Q}_\mu}{R^3} + \dots \quad (4)$$

with $u_\alpha = R_\alpha/R$; $\mathbf{p}_\mu \mathbf{u} = p_\mu^\alpha u_\alpha$, $\mathbf{u} : \hat{Q}_\mu : \mathbf{u} = Q_\mu^{\alpha\beta} u_\alpha u_\beta$, $Sp\hat{Q}_\mu = Q_\mu^{\alpha\alpha}$ and $\mu = (x, y)$. This is the most general expression for the director field. It differs from usual electrodynamics with $-Sp\hat{Q}_\mu/R^3$ term. We shall show below that it is not zero when the axial symmetry breaking takes place. We note that the multiple expression does not depend on the anchoring strength. It is valid on far distances for any anchoring, weak and strong, without topological defects or with them. Of course in order to find multiple coefficients we need to solve the problem in the near nonlinear area either with computer simulation or Ansatz functions. Let's imagine that we have found all multiple coefficients for the particular particle (for instance with help of computer simulation): 20 different values, with 10 for every n_μ . So for every $\mu = (x, y)$ we have one elastic charge q_μ , three components of the vector dipole moment p_μ^α and six components of the quadrupole tensor $\hat{Q}_\mu$ (it is symmetric $Q_\mu^{\alpha\beta} = Q_\mu^{\beta\alpha}$, but $Sp\hat{Q}_\mu \neq 0$ in general). How particles will elastically interact if we know all these 20 multiple coefficients. We want to obtain analytical interaction potential. Our task is to build effective harmonic free energy functional which includes all multiple coefficients and nothing else. It must be invariant and it's Euler-Lagrange equations for one particle must have solution in the form of multiple expansion (4). It is easy to check that such functional has the following form:

$$F = K \int d^3x \left\{ \frac{(\nabla n_\mu)^2}{2} - 4\pi q_\mu(\mathbf{x}) n_\mu - 4\pi p_\mu^\alpha(\mathbf{x}) \partial_\alpha n_\mu - 4\pi Q_\mu^{\alpha\beta}(\mathbf{x}) \partial_\alpha \partial_\beta n_\mu \right\} \quad (5)$$

In the case of N particles $q_\mu(\mathbf{x}) = \sum_i q_{\mu,i} \delta(\mathbf{x} - \mathbf{x}_i)$; $p_\mu^\alpha(\mathbf{x}) = \sum_i p_{\mu,i}^\alpha \delta(\mathbf{x} - \mathbf{x}_i)$; $Q_\mu^{\alpha\beta}(\mathbf{x}) = \sum_i Q_{\mu,i}^{\alpha\beta} \delta(\mathbf{x} - \mathbf{x}_i)$ with i being the number of the particle. Euler-Lagrange equations for functional (5) are Poisson like:

$$\Delta n_\mu = -4\pi q_\mu(\mathbf{x}) + 4\pi [\partial_\alpha (p_\mu^\alpha(\mathbf{x})) - \partial_\alpha \partial_\beta Q_\mu^{\alpha\beta}(\mathbf{x})] \quad (6)$$

Substitution of this equation into (5) leads to the following expression for the total energy of the system: particles and liquid crystal:

$$F_{total} = -2\pi K \int d^3x \left\{ q_\mu(\mathbf{x}) n_\mu^{ex} + (\mathbf{p}_\mu(\mathbf{x}) \cdot \nabla) n_\mu^{ex} + (\hat{Q}_\mu(\mathbf{x}) : \nabla : \nabla) n_\mu^{ex} \right\} \quad (7)$$

where n_μ^{ex} -exact solution of the (6). For unlimited space the solution of equation (6) has standard expression:

$$n_\mu(\mathbf{x}) = \int d^3\mathbf{x}' \frac{1}{|\mathbf{x} - \mathbf{x}'|} [q_\mu(\mathbf{x}') - \partial'_\alpha (p_\mu^\alpha(\mathbf{x}')) + \partial'_\alpha \partial'_\beta Q_\mu^{\alpha\beta}(\mathbf{x}')] \quad (8)$$

For one particle we have $q_\mu(\mathbf{x}) = q_\mu \delta(\mathbf{x})$, $p_\mu^\alpha(\mathbf{x}) = p_\mu^\alpha \delta(\mathbf{x})$, $Q_\mu^{\alpha\beta}(\mathbf{x}) = Q_\mu^{\alpha\beta} \delta(\mathbf{x})$ which leads us immediately to the general expression (4). For axially symmetric particles $q_\mu = 0$ and the director take the form $n_x = p \frac{x}{R^3} + 6Q \frac{xz}{R^5}$, $n_y = p \frac{y}{R^3} + 6Q \frac{yz}{R^5}$. In Lubensky notation [7] $Q = \frac{c}{3}$.

Actually in the construction of the functional (5) we follow Lubensky ideology but with one significant difference. In the paper [7] authors tried to build invariants of vector $\mathbf{P}$, tensor C_{ij} and director vector $\mathbf{n}$, which has three components, not two. So obtained invariants are of the type $-\mathbf{P} \cdot \mathbf{n}(\nabla \mathbf{n})$, $(\nabla \mathbf{n}) \mathbf{n} \cdot \nabla (n_i C_{ij} n_j)$ and include nonlinear terms of n_μ . In linear approximation authors find the following main elastic functional:

$$F = K \int d^3x \left\{ \frac{(\nabla n_\mu)^2}{2} - 4\pi P_z(\mathbf{x}) \partial_\mu n_\mu - 4\pi C_{zz}(\mathbf{x}) \partial_z \partial_\mu n_\mu \right\} \quad (9)$$

This functional is axially symmetric, it correctly describes axially symmetric particles. But in the case of axial symmetry breaking it does not valid. In the case of axial symmetry breaking we need to take into account coulomb like expression and to introduce two different vectors $\mathbf{p}_x, \mathbf{p}_y$ and two tensors $\hat{Q}_x, \hat{Q}_y$ for each component μ . Using them we find new functional (5) which describes system of any particles immersed into NLC. Let's show how (5) proceeds to the (9) for axially symmetric particles. In this case $\mathbf{p}_x = (p, 0, 0)$; $\mathbf{p}_y = (0, p, 0)$ and quadrupole tensors are of the form:

$$\hat{Q}_x = \begin{pmatrix} 0 & 0 & Q \\ 0 & 0 & 0 \\ Q & 0 & 0 \end{pmatrix} \hat{Q}_y = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & Q \\ 0 & Q & 0 \end{pmatrix}. \quad (10)$$

Then automatically (5) proceeds to the (9) with $P_z(\mathbf{x}) = p(\mathbf{x})$; $C_{zz}(\mathbf{x}) = 2Q(\mathbf{x})$.

It is necessary to note that functional (5) defines energy of the colloidal particle in the director deformations created by any external sources. So if we have some deformed NLC the particle will move in the way to minimize (5).

Director field from N immersed particles will be superposition of deformations (4) from all particles $n_\mu(\mathbf{x}) = \sum_i \frac{q_{\mu,i}}{|\mathbf{x}-\mathbf{x}_i|} + \sum_i \frac{\mathbf{p}_{\mu,i}(\mathbf{x}-\mathbf{x}_i)}{|\mathbf{x}-\mathbf{x}_i|^3} + \sum_i \frac{3(\mathbf{x}-\mathbf{x}_i):\hat{Q}_{\mu,i}:(\mathbf{x}-\mathbf{x}_i)}{|\mathbf{x}-\mathbf{x}_i|^5} - \sum_i \frac{Sp\hat{Q}_{\mu,i}}{|\mathbf{x}-\mathbf{x}_i|^3}$. Substituting this director into (7) and neglecting with self energy divergent parts we come to the fact that total energy of the system may be presented as the sum of pair interactions:

$$F_{total} = \sum_i \sum_{j>i} U^{ij}(\mathbf{x}_i - \mathbf{x}_j) \quad (11)$$

$$U^{ij} = U_{qq} + U_{qd} + U_{dd} + U_{qQ} + U_{dQ} + U_{QQ} \quad (12)$$

$$U_{qq} = -4\pi K \frac{q_{\mu,i} q_{\mu,j}}{r_{ij}^2} \quad (13)$$

$$U_{qd} = -4\pi K \frac{q_{\mu,i}(\mathbf{p}_{\mu,j} \cdot \mathbf{u}_{ij}) - q_{\mu,j}(\mathbf{p}_{\mu,i} \cdot \mathbf{u}_{ij})}{r_{ij}^2} \quad (14)$$

$$U_{dd} = -4\pi K \frac{(\mathbf{p}_{\mu,i} \cdot \mathbf{p}_{\mu,j}) - 3(\mathbf{p}_{\mu,i} \cdot \mathbf{u}_{ij})(\mathbf{p}_{\mu,j} \cdot \mathbf{u}_{ij})}{r_{ij}^3} \quad (15)$$

$$U_{qQ} = -12\pi K \left\{ \frac{q_{\mu,i}(\mathbf{u}_{ij} : \hat{Q}_{\mu,j} : \mathbf{u}_{ij}) + q_{\mu,j}(\mathbf{u}_{ij} : \hat{Q}_{\mu,i} : \mathbf{u}_{ij})}{r_{ij}^3} - \frac{(q_{\mu,i} Sp\hat{Q}_{\mu,j} + q_{\mu,j} Sp\hat{Q}_{\mu,i})}{3r_{ij}^3} \right\} \quad (16)$$

$$U_{dQ} = -12\pi K \left\{ \frac{(\mathbf{p}_{\mu,i} \cdot \mathbf{u}_{ij}) Sp\hat{Q}_{\mu,j} - (\mathbf{p}_{\mu,j} \cdot \mathbf{u}_{ij}) Sp\hat{Q}_{\mu,i}}{r_{ij}^4} + 2 \frac{(\mathbf{p}_{\mu,i} : \hat{Q}_{\mu,j} : \mathbf{u}_{ij} - \mathbf{p}_{\mu,j} : \hat{Q}_{\mu,i} : \mathbf{u}_{ij})}{r_{ij}^4} \right\} - \quad (17)$$

$$-60\pi K \frac{(\mathbf{u}_{ij} : \hat{Q}_{\mu,i} : \mathbf{u}_{ij})(\mathbf{p}_{\mu,j} \cdot \mathbf{u}_{ij}) - (\mathbf{u}_{ij} : \hat{Q}_{\mu,j} : \mathbf{u}_{ij})(\mathbf{p}_{\mu,i} \cdot \mathbf{u}_{ij})}{r_{ij}^4}$$

$$U_{QQ} = -12\pi K \left\{ \frac{2\hat{Q}_{\mu,i} :: \hat{Q}_{\mu,j} - 20\mathbf{u}_{ij} : \hat{Q}_{\mu,i} : \hat{Q}_{\mu,j} : \mathbf{u}_{ij} + 35(\mathbf{u}_{ij} : \hat{Q}_{\mu,i} : \mathbf{u}_{ij})(\mathbf{u}_{ij} : \hat{Q}_{\mu,j} : \mathbf{u}_{ij})}{r_{ij}^5} \right\} - \quad (18)$$

$$-12\pi K \frac{Sp\hat{Q}_{\mu,i} Sp\hat{Q}_{\mu,j}}{r_{ij}^5}$$

where $\mathbf{u}_{ij} = \frac{\mathbf{x}_i - \mathbf{x}_j}{|\mathbf{x}_i - \mathbf{x}_j|}$ and $r_{ij} = |\mathbf{x}_i - \mathbf{x}_j|$. These expressions are the most general elastic pair interaction potentials between ordinary particles immersed into NLC. They are valid for particles with topological defects and without them, for any type and strength of the anchoring coefficient. Coulomb-like elastic attraction and repulsion between inclined ellipsoidal particles was found in [16],[17] with help of computer simulation. We consider that Coulomb-like elastic interaction takes place between any non-symmetrical particles with broken axial symmetry (for instance non-symmetrical triangles or cubes with different anchoring on sides) even in the case of zero torque applied to the particles.

If particles are the same then $U_{qd} \equiv 0, U_{dQ} \equiv 0$. We see that formulas (16)-(18) contains new terms connected with $Sp\hat{Q}_\mu$ which are absent in the classical electrostatics. Let's analyze this result in comparison with electrostatics. In electrostatics there is the charge density distribution $\rho(\mathbf{x})$ in the Poisson equation $\Delta\varphi = -4\pi\rho$ and linear electrostatics works *inside* this density. So *inside* $\rho(\mathbf{x})$ there is superposition principle. But in nematostatics there is *principle* difference on small distances - the superposition principle does not work because of nonlinear deformations. So in case of strong anchoring we consider that the notion of elastic charge density is unreasonable. That's why in nematostatics we introduce the concept of elastic charge q_μ , dipole moment $\mathbf{p}_\mu$, quadrupole moment $Q_\mu^{\alpha\beta}$ not as compound expressions like in electrostatics (full charge is equal the sum of compound charges $q_{total} = \sum_i q_i$, dipole moment as $\mathbf{P} = \sum_i q_i \mathbf{r}_i$ and so on), but as primary quantities for each particle, the quantities which are just the linear asymptotics of exact *nonlinear solutions*.

As well in nemato-statics surface effects can make $Sp\hat{Q}_\mu \neq 0$. Let's show this on the example of small particles with weak anchoring without topological defects. Then deformations are small anywhere $\mathbf{n}(\mathbf{R}) \approx (n_x, n_y, 1)$, $|n_\mu| \ll 1$ ($\mu = x, y$) and surface energy takes the form $\oint dS W(\nu \cdot \mathbf{n})^2 \approx const + 2W \oint dS \nu_\mu \nu_3 n_\mu(\mathbf{s})$. Taylor expansion of the director concerning center of the particle brings: $n_\mu(s) = n_\mu(x_0) + (\mathbf{l} \cdot \nabla) n_\mu(x_0) + \frac{1}{2}(\mathbf{l} \cdot \nabla)^2 n_\mu(x_0)$ where $\mathbf{l}$ is the vector from the center $\mathbf{x}_0$ to the point on the surface $\mathbf{s} = \mathbf{x}_0 + \mathbf{l}$. Then we automatically obtain that free energy (1) takes the form (5) with such multiple coefficients: $\tilde{q}_\mu = -\frac{W}{2\pi K} \oint dS \nu_3 \nu_\mu$, $\tilde{\mathbf{p}}_\mu = -\frac{W}{2\pi K} \oint dS \nu_3 \nu_\mu \mathbf{l}(s)$, $\tilde{Q}_\mu^{\alpha\beta} = -\frac{W}{4\pi K} \oint dS \nu_3 \nu_\mu l_\alpha(s) l_\beta(s)$. These quantities are of course approximate as Taylor extension is not exact but we consider that qualitative estimates may be correctly performed. Let's note that substituting them into U_{ij} we naturally come to the results of papers [9, 10].

So we may clearly see that $Sp\tilde{Q}_\mu = -\frac{W}{4\pi K} \oint dS \nu_3 \nu_\mu l^2(s) \approx \langle l^2 \rangle \tilde{q}_\mu \neq 0(s)$ if $\tilde{q}_\mu \neq 0$, so that $Sp\tilde{Q}_\mu$ and $\tilde{q}_\mu$ are approximately proportional to each other (here $\langle l^2 \rangle$ is average square of the distance from the surface to the center of the particle). As the elastic charge occurs when axial symmetry is broken [10] we come to the conclusion that in that case as well $Sp\tilde{Q}_\mu \neq 0$. In the paper [17] authors have found Coulomb-like elastic interaction between two inclined ellipsoidal particles using computer simulation. As well there analytical expression for the elastic charge for the case of small anchoring was obtained using $\tilde{q}_\mu = -\frac{W}{2\pi K} \oint dS \nu_3 \nu_\mu$. Then it was found that $\tilde{q}_x = -\frac{W}{2\pi K} F(b, \varepsilon) \cos \psi \sin \theta \cos \theta$, $\tilde{q}_y = -\frac{W}{2\pi K} F(b, \varepsilon) \sin \psi \sin \theta \cos \theta$, where θ is the angle between long axis and $\mathbf{n}_0$, ψ is the angle between the horizontal projection of the particle and direction of axis x , b is the length of the long axis and ε is the eccentricity $\varepsilon = \sqrt{1 - c^2/b^2}$ with c being the short axis of the ellipsoid and:

$$F(b, \varepsilon) = \pi b^2 \sqrt{1 - \varepsilon^2} \left\{ \frac{\sqrt{1 - \varepsilon^2}}{\varepsilon^2} (2\varepsilon^2 - 3) + \frac{\arcsin \varepsilon}{\varepsilon^3} (3 - 4\varepsilon^4) \right\}. \quad (19)$$

In the case of small eccentricities it has the asymptotic expansion $F(b, \varepsilon)_{\varepsilon \rightarrow 0} = -\frac{10}{6} \pi b^2 \varepsilon^2 + o(\varepsilon^2)$. Here we come to an interesting conclusion, that the elastic charge goes to zero as square of the eccentricity $\tilde{q}_\mu \propto \varepsilon^2$. Then we may estimate $Sp\tilde{Q}_\mu \approx \tilde{q}_\mu \langle l^2 \rangle$, where $\langle l^2 \rangle$ is something average between b^2 and c^2 .

In this paper the new free energy functional which describes elastic interaction between arbitrary colloidal particles and NLC has been proposed. It contains multiple coefficients of each colloidal particle and proceeds to Lubensky functional for axially symmetric particles. This approach enables to find elastic interaction energy between emerged particles in the most general case for arbitrary anchoring strength on the surface. The procedure to find interaction energy consists of two steps - first, multiple coefficients for director field are found using different methods like computer simulation, variational ansatz and so on, second - these coefficients are used in the obtained analytical expressions for elastic potentials. It was first time shown that spur of the quadrupole moment tensor is different from zero $Sp\hat{Q}_\mu \neq 0$ in the case of axial symmetry breaking of the director field around colloidal particle in contrast to classical electrostatics.

- [2] P.Poulin and D.A.Weitz, Phys.Rev. E **57**, 626 (1998).
- [3] I. Mušević, M. Škarabot, U.Tkalec, M.Ravnik and S.Žumer Science **313**, 954, (2006).
- [4] V.Nazarenko, A.Nych and B.Lev, Phys.Rev.Lett , **87**,075504 (2001).
- [5] I. I. Smalyukh, S. Chernyshuk, B. I. Lev, A. B. Nych, U. Ognysta, V.G. Nazarenko, and O. D. Lavrentovich, Phys. Rev. Lett. **93**, 117801, (2004).
- [6] K.Takahashi, M. Ichikawa and Y.Kimura, Phys. Rev. E., **77**, 020703(R),(2008)
- [7] T.C.Lubensky, D.Pettey, N.Currier and H.Stark, Phys.Rev.E **57**, 610 (1998).
- [8] S.L.Lopatnikov and V.A.Namiot, Zh. Eksp. Teor. Fiz. **75**, 361 (1978) [Sov. Phys. JETP **48**, 180 (1978)].
- [9] B.I.Lev and P.M.Tomchuk, Phys.Rev.E **59**, 591 (1999).
- [10] B.I.Lev,S.B.Chernyshuk,P.M.Tomchuk and H.Yokoyama, Phys.Rev.E **65**,021603,(2002)
- [11] V. M. Pergamenshchik and V. A. Uzunova, Eur. Phys. J. E , **23**,161 (2007)
- [12] V. M. Pergamenshchik and V. A. Uzunova, Phys. Rev. E **76**,011707 (2007)
- [13] Jun-Ici Fukuda and Slobodan Zumer, Phys. Rev. E., **79**, 041703,(2009)
- [14] V.A.Raghunathan et.al, Mol.Cryst.Liq.Cryst. **288**, 181 (1996).
- [15] F.Brochard and P.G. De Gennes, J.Phys.(Paris)
- [16] M.Ravnik, PhD thesis, Ljubljana, Slovenia, 2009
- [17] S.Chernyshuk, M.Ravnik, B.Lev, S.Žumer, to be published.